

1. Solve each of the following.

a)  $\frac{7}{2x-1} \leq -3$     b)  $\frac{x-1}{3x+1} \leq \frac{1}{3}$     c)  $\frac{x}{x+4} \geq \frac{2}{3}$     d)  $\frac{x+3}{x-3} < 2$

2. We draw tangent lines to a circle from a point  $P$  outside of the circle. The line segment between  $P$  and a point of tangency is 3 units long. The line segment connecting  $P$  and the center of the circle intersects the circle in point  $Q$ . Line segment  $PQ$  is  $\sqrt{3}$  units long. Compute the angle formed between the two tangent lines.

3. Compute the exact value of each of the following. Rationalize denominators and simplify your answer.

a)  $\cos 15^\circ$     b)  $\sin 15^\circ$     c)  $\tan 75^\circ$

4. Find an equation for all tangent lines drawn from  $P(2, 13)$  to the graph of  $y = -\frac{1}{2}x^2 + 3x + 1$ .

5. Prove each of the following identities.

a)  $\cot 2x = \frac{\cot^2 x - 1}{2 \cot x}$

d)  $\sin\left(\frac{\pi}{2} - x\right) = \cos x$

b)  $4 \sin^4 x = 1 - 2 \cos 2x + \cos^2 2x$

e)  $\frac{\tan(45^\circ - x)}{\tan(45^\circ + x)} = \frac{1 - \sin 2x}{1 + \sin 2x}$

c)  $\cos 3x = 4 \cos^3 x - 3 \cos x$

f)  $\sin(30^\circ + \beta) + \sin(30^\circ - \beta) = \cos \beta$

6. Simplify completely or re-write each of the following as a single logarithm.

a)  $\log_3 8 - \log_9 24$

b)  $\log_2 3 - \log_4 6$

c)  $(\log_2 3)(\log_3 5)(\log_5 8)$

7. Suppose that  $\tan x = -\frac{4}{5}$ . Compute the exact value of  $\sin 2x$ .

8. Given  $A(5, 2)$  and  $B(-11, 6)$ , find an equation for the perpendicular bisector of  $A$  and  $B$ .

9. Simplify each of the following. Use exact values.

a)  $\frac{\sin\left(\frac{7\pi}{2}\right) \cos\left(-\frac{11\pi}{6}\right)}{\tan\left(\frac{13\pi}{3}\right)}$

b)  $-\cos(12^\circ) \sin(72^\circ) + \sin(12^\circ) \cos(72^\circ)$

10. Suppose that  $a = \log_2 5$  and  $b = \log_3 6$ . Write  $\log_{12} 60$  in terms of  $a$  and  $b$ .

11. Simplify each of the following. (Write it in terms of  $\sin \alpha$ ,  $\cos \alpha$ , and/or  $\tan \alpha$ .)

a)  $\sin(\alpha + 90^\circ)$

b)  $\cos(\alpha + 90^\circ)$

c)  $\tan(\alpha - 180^\circ)$

d)  $\cos(\alpha - 180^\circ)$

12. Which of the following functions are not one-to-one?

a)  $f(x) = -\ln x$     b)  $g(x) = \frac{1}{3}x + 8$     c)  $h(x) = x^3 - x$     d)  $f(x) = \frac{3x-8}{x+3}$

13. Simplify each of the following.

a)  $\log_3(\tan 60^\circ) + \log_2(\sin 45^\circ) - \ln(\tan 45^\circ)$

c)  $\frac{\tan 160^\circ - \tan 25^\circ}{1 + \tan 160^\circ \tan 25^\circ}$

b)  $\sec 330^\circ - \csc(-315^\circ) + \cot 60^\circ$

d)  $\frac{\tan 15^\circ + 1}{\tan 15^\circ - 1}$

14. Find the domain of  $f(x) = \frac{x^2 - 4}{5 - \sqrt{2x - 4}}$

15. Prove each of the following co-function identities.

$$\text{a) } \sin\left(\frac{\pi}{2} - \alpha\right) = \cos \alpha \quad \text{b) } \cos\left(\frac{\pi}{2} - \alpha\right) = \sin \alpha \quad \text{c) } \tan\left(\frac{\pi}{2} - \alpha\right) = \cot \alpha$$

16. Suppose that  $\alpha$  and  $\beta$  are acute angles with  $\sin \alpha = \frac{1}{3}$  and  $\cos \beta = \frac{5}{13}$ . Compute the exact value of each of the following.

$$\begin{array}{llll} \text{a) } \cos \alpha & \text{d) } \tan \beta & \text{g) } \sin(\alpha + \beta) & \text{j) } \cos(\alpha - \beta) \\ \text{b) } \tan \alpha & \text{e) } \sin 2\alpha & \text{h) } \sin(\alpha - \beta) & \text{k) } \tan 2\alpha \\ \text{c) } \sin \beta & \text{f) } \cos 2\beta & \text{i) } \cos(\alpha + \beta) & \text{l) } \tan 2\beta \end{array}$$

17. Suppose that  $\sin A = -\frac{3}{5}$ . Compute the exact value of each of the following.

$$\text{a) } \sin 2A \quad \text{b) } \cos 2A$$

18. Suppose that  $\sin B = \frac{1}{3}$  and  $B$  is not in the first quadrant. Compute the exact value of each of the following.

$$\text{a) } \sec B \quad \text{b) } \tan 2B \quad \text{c) } \cos 2B \quad \text{d) } \tan\left(B - \frac{\pi}{4}\right)$$

19. The expression  $\frac{2 \sin x}{\cos x - \sin x \tan x}$  is equivalent to which of the following?

$$\text{A) } \tan 2x \quad \text{B) } \cot 2x \quad \text{C) } \tan x \quad \text{D) } \cot x \quad \text{E) } \sec x$$

20. Compute the sum  $83 + 86 + 89 + \dots + 203$ .

21. Compute the exact value of  $\frac{\tan\left(\frac{4\pi}{3}\right) - \tan\left(\frac{\pi}{12}\right)}{1 + \tan\left(\frac{4\pi}{3}\right)\tan\left(\frac{\pi}{12}\right)}$ . (Hint: there is an easy way and also a difficult way to do this.)

22. Solve each of the following equations.

$$\begin{array}{lll} \text{a) } \tan^2 x + \tan x = 0 & \text{c) } \sin x = \sin 2x & \text{e) } \cos x - 2 \sin^2 x = 1 \\ \text{b) } \sin^2 x = \frac{1}{2} & \text{d) } \sin x + 2 \cos^2 x = 1 & \end{array}$$

23. Find the domain for each of the following functions.

$$\text{a) } f(x) = \log_2(x^2 - 9) \quad \text{b) } \log_2(x + 3) + \log_2(x - 3) \quad \text{c) } \ln\left(\frac{x + 1}{x - 1}\right) \quad \text{d) } \frac{\ln(x + 1)}{\ln(x - 1)}$$

24. Circle  $C_1$  has a radius 5 unit long. Circle  $C_2$  has a radius 11 unit long. The centers are at a distance of 12 units from each other. We draw the lines tangent to both circles.

- Find an approximation of the angle formed by the two tangent lines.
- Compute the distance between the two points of tangency on one of the common tangent lines.

25. Compute the exact value of  $\tan \alpha$  if we know that  $\tan 2\alpha = \frac{8}{15}$ .

26. A lattice point is a point on the coordinate system of whose both coordinates are integers. Can you find the equation of a line with

- No lattice points?
- Exactly one lattice point?
- Exactly two lattice points?

27. We leave \$1 in a bank that promises an annual compound interest rate of 5%. We notify the bank that a relative will withdraw all money and close the account once it reached \$1000 000. How long will the bank have to maintain the account?
28. Consider the triangle determined by the points  $A(-1, 2)$ ,  $B(6, 8)$ , and  $C(7, 6)$ . Compute the exact value of the area of the triangle.

## Answers

1. a)  $\left[-\frac{2}{3}, \frac{1}{2}\right)$    b)  $\left(-\frac{1}{3}, \infty\right)$    c)  $(-\infty, -4) \cup [8, \infty)$    d)  $(-\infty, 3) \cup (9, \infty)$

2.  $60^\circ$    3. a)  $\frac{\sqrt{6} + \sqrt{2}}{4}$    b)  $\frac{\sqrt{6} - \sqrt{2}}{4}$    c)  $2 + \sqrt{3}$    4.  $y = 5x + 3$  and  $y = -3x + 19$

5. a)  $\cot 2x = \frac{\cot^2 x - 1}{2 \cot x}$

$$\begin{aligned} \text{RHS} &= \frac{\cot^2 x - 1}{2 \cot x} = \frac{\frac{\cos^2 x}{\sin^2 x} - 1}{2 \frac{\cos x}{\sin x}} = \frac{\frac{\cos^2 x - \sin^2 x}{\sin^2 x}}{2 \frac{\cos x}{\sin x}} = \frac{\frac{\cos^2 x - \sin^2 x}{\sin^2 x}}{\frac{2 \cos x}{\sin x}} = \frac{\cos^2 x - \sin^2 x}{\sin^2 x} \cdot \frac{\sin x}{2 \cos x} \\ &= \frac{\cos^2 x - \sin^2 x}{2 \sin x \cos x} = \frac{\cos 2x}{\sin 2x} = \cot 2x = \text{LHS} \end{aligned}$$

b)  $4 \sin^4 x = 1 - 2 \cos 2x + \cos^2 2x$

$$\begin{aligned} \text{LHS} &= 4 \sin^4 x = 4 (\sin^2 x)^2 = (2 \sin^2 x)^2 = (2 \sin^2 x - 1 + 1)^2 = (-\cos 2x + 1)^2 = \\ &= \cos^2 2x - 2 \cos 2x + 1 = \text{RHS} \end{aligned}$$

c)  $\cos 3x = 4 \cos^3 x - 3 \cos x$

$$\begin{aligned} \text{LHS} &= \cos 3x = \cos(x + 2x) = \cos x \cos 2x - \sin x \sin 2x = \cos x (2 \cos^2 x - 1) - \sin x (2 \sin x \cos x) = \\ &= 2 \cos^3 x - \cos x - 2 \sin^2 x \cos x = 2 \cos^3 x - \cos x - 2(1 - \cos^2 x) \cos x = \\ &= 2 \cos^3 x - \cos x - 2 \cos x + 2 \cos^3 x = 4 \cos^3 x - 3 \cos x = \text{RHS} \end{aligned}$$

d)  $\sin\left(\frac{\pi}{2} - x\right) = \cos x$

$$\text{LHS} = \sin\left(\frac{\pi}{2}\right) \cos x - \cos\left(\frac{\pi}{2}\right) \sin x = 1 \cos x - 0 \sin x = \cos x = \text{RHS}$$

e)  $\frac{\tan(45^\circ - x)}{\tan(45^\circ + x)} = \frac{1 - \sin 2x}{1 + \sin 2x}$

$$\begin{aligned} \text{LHS} &= \frac{\tan(45^\circ - x)}{\tan(45^\circ + x)} = \frac{\frac{\tan 45^\circ - \tan x}{1 + \tan 45^\circ \tan x}}{\frac{\tan 45^\circ + \tan x}{1 - \tan 45^\circ \tan x}} = \frac{\frac{1 - \tan x}{1 + 1 \cdot \tan x}}{\frac{1 + \tan x}{1 - 1 \cdot \tan x}} = \frac{1 - \tan x}{1 + \tan x} \cdot \frac{1 - \tan x}{1 + \tan x} = \left(\frac{1 - \frac{\sin x}{\cos x}}{1 + \frac{\sin x}{\cos x}}\right)^2 \\ &= \frac{\left(1 - \frac{\sin x}{\cos x}\right)^2}{\left(1 + \frac{\sin x}{\cos x}\right)^2} = \frac{\left(1 - \frac{\sin x}{\cos x}\right)^2 \cdot \cos^2 x}{\left(1 + \frac{\sin x}{\cos x}\right)^2 \cdot \cos^2 x} = \frac{\left(\left(1 - \frac{\sin x}{\cos x}\right) \cos x\right)^2}{\left(\left(1 + \frac{\sin x}{\cos x}\right) \cos x\right)^2} = \frac{(\cos x - \sin x)^2}{(\cos x + \sin x)^2} \\ &= \frac{\cos^2 x + \sin^2 x - 2 \sin x \cos x}{\cos^2 x + \sin^2 x + 2 \sin x \cos x} = \frac{1 - \sin 2x}{1 + \sin 2x} = \text{RHS} \end{aligned}$$

f)  $\sin(30^\circ + \beta) + \sin(30^\circ - \beta) = \cos \beta$

$$\begin{aligned} \text{LHS} &= \sin(30^\circ + \beta) + \sin(30^\circ - \beta) = \sin 30^\circ \cos \beta + \cos 30^\circ \sin \beta + \sin 30^\circ \cos \beta - \cos 30^\circ \sin \beta \\ &= 2 \sin 30^\circ \cos \beta = 2 \cdot \frac{1}{2} \cos \beta = \cos \beta = \text{RHS} \end{aligned}$$

6. a)  $\log_3 \left( \frac{2\sqrt{6}}{3} \right)$  or  $\log_3(2\sqrt{6}) - 1$     b)  $\log_2 \left( \frac{\sqrt{6}}{2} \right)$  or  $\log_2(\sqrt{6}) - 1$     c) 3    7.  $-\frac{40}{41}$     8.  $y = 4x + 16$

9. a)  $-\frac{1}{2}$     b)  $-\frac{\sqrt{3}}{2}$     10.  $\frac{-a+b+ab}{2b-1}$     11. a)  $\cos \alpha$     b)  $-\sin \alpha$     c)  $\tan \alpha$     d)  $-\cos \alpha$

12. only  $h$  is not one-to-one    13. a) 0    b)  $\sqrt{3} - \sqrt{2}$     c)  $-1$     d)  $-\sqrt{3}$

14.  $\{x \in \mathbb{R} : x \geq 2 \text{ and } x \neq 14.5\}$  in interval notation:  $[2, 14.5) \cup (14.5, \infty)$

15. a)  $\sin \left( \frac{\pi}{2} - \alpha \right) = \cos \alpha$

$$\sin \left( \frac{\pi}{2} - \alpha \right) = \sin \frac{\pi}{2} \cos \alpha - \cos \frac{\pi}{2} \sin \alpha = 1 \cdot \cos \alpha - 0 \cdot \sin \alpha = \cos \alpha$$

b)  $\cos \left( \frac{\pi}{2} - \alpha \right) = \sin \alpha$

$$\cos \left( \frac{\pi}{2} - \alpha \right) = \cos \frac{\pi}{2} \cos \alpha + \sin \frac{\pi}{2} \sin \alpha = 0 \cdot \cos \alpha + 1 \cdot \sin \alpha = \sin \alpha$$

c)  $\tan \left( \frac{\pi}{2} - \alpha \right) = \cot \alpha$  We can not use the difference formula for tangent here, because  $\tan \frac{\pi}{2}$  is undefined. Instead, we need to use a compound angle formula separately for sine and cosine (see parts a and b).

$$\tan \left( \frac{\pi}{2} - \alpha \right) = \frac{\sin \left( \frac{\pi}{2} - \alpha \right)}{\cos \left( \frac{\pi}{2} - \alpha \right)} = \frac{\cos \alpha}{\sin \alpha} = \cot \alpha$$

16. a)  $\frac{2\sqrt{2}}{3}$     b)  $\frac{\sqrt{2}}{4}$     c)  $\frac{12}{13}$     d)  $\frac{12}{5}$     e)  $\frac{4\sqrt{2}}{9}$     f)  $-\frac{119}{169}$     g)  $\frac{5+24\sqrt{2}}{39}$     h)  $\frac{5+24\sqrt{2}}{39}$   
 i)  $\frac{10\sqrt{2}-12}{39}$     j)  $\frac{10\sqrt{2}+12}{39}$     k)  $\frac{4\sqrt{2}}{7}$     l)  $-\frac{120}{119}$

17. a)  $\pm \frac{24}{25}$     b)  $\frac{7}{25}$     18. a)  $-\frac{3\sqrt{2}}{4}$     b)  $-\frac{4\sqrt{2}}{7}$     c)  $\frac{8}{9}$     d)  $-\frac{1}{2}$     19. A    20. 5863    21. 1

22. a)  $-\frac{\pi}{4} + k\pi$  and  $k\pi$  where  $k \in \mathbb{Z}$     b)  $\frac{\pi}{4} + \frac{k\pi}{2}$  where  $k \in \mathbb{Z}$     c)  $k\pi$  or  $\pm \frac{\pi}{3} + 2k\pi$  where  $k \in \mathbb{Z}$   
 d)  $x = \frac{\pi}{2} + 2k\pi$  or  $x = -\frac{\pi}{6} + 2k\pi$  or  $x = -\frac{5\pi}{6} + 2k\pi$  where  $k \in \mathbb{Z}$     e)  $x = 2k\pi$  where  $k \in \mathbb{Z}$

23. a)  $(-\infty, -3) \cup (3, \infty)$     b)  $(3, \infty)$     c)  $(-\infty, -1) \cup (1, \infty)$     d)  $(1, 2) \cup (2, \infty)$

24. a)  $60^\circ$     b)  $\sqrt{108}$  unit =  $6\sqrt{3}$  unit    25.  $-4$  or  $\frac{1}{4}$

26. a)  $y = \frac{5}{3}$     b)  $y = \sqrt{2}x$  will only contain the origin    c) there isn't such a line

27. about 284 years    28.  $10 \text{ unit}^2$